



2019

2^{ème} Conférence Intensification Durable

Session 1.2 Agroforestry

Vertebrates contribute to natural control of the millet head miner in tree-crop agroforestry systems

8—10 october 2019

Hotel Ngor Diarama
Dakar, Sénégal

Ahmadou Sow, Djiby Seye, Emile Faye,
Laure Benoit, Maxime Galan, Julien Haran,
Thierry Brévault



Presentation plan

2

- Biodiversity services in agroecosystems
- Tree-crop agroforestry systems promote natural pest control
- On the role of birds and bats as insect predators
- Take-home message



Crop pests

3

A major obstacle to the increase of agricultural production



What about natural pest control?

4

Intensification of agriculture

Simplification of agricultural landscapes



Loss of biodiversity and associated ecosystem services
such as natural pest control




Cultivated ecosystems more
susceptible to pests

Trees can promote natural pest control

5

By providing habitats to natural enemies in agroecosystems



- Refuges
- Food resources
- Alternative preys
- Microclimate
- Barrier to pest colonization
- Etc.

However, trees can also benefit pests...

What did meta-analyses say?

6

Agroforestry practices generally result in...

**Natural
enemies**



Pests



in perennial crops
(coffee, cocoa and plantain)



in annual crops
(e.g. maize, rice and beans)

(Pumariño et al. 2015)



(Ogol et al. 1999)

Tree-crop systems as a case study

7

Traditional agroforestry parkland systems in Senegal



Pearl millet as a staple crop


8



Wanted: The millet head miner

9

Heliocheilus albipunctella (Lepidoptera, Noctuidae)




- Monophagous, only one generation a year
- 30-95% infested panicles - 2-20% grain yield loss (Sow et al. 2017)

Tree-crop agroforestry systems promote natural pest control in millet fields

10

- Natural pest control increases with the abundance of trees in the landscape




(Soti et al. 2019)

Food webs and natural pest control

11

- High throughput molecular tools (metabarcoding) were used to document food web structures (parasitism and predation)




- What about vertebrates?

(Sow et al. 2018, 2019)

Case study:

Trees, vertebrates and natural pest control

12




10 Sites : high vs low tree density in a 1000 m-buffer (landscape)

2 fields per site : open vs closed in a 100 m-buffer (local)

Case study:

Trees, vertebrates and natural pest control

13



In each site, trees were geolocated and identified at the species level using drone imagery and remote sensing.




Material and methods

14


1- Exclusion of natural enemies

2- Field monitoring of birds




Test of research hypothesis

15




Abundance and diversity of tree cover

16



Insectivorous birds and bats


17



Taphozous mauritanus

Detection of pest DNA in feces

18



Ploceus cucullatus




Taphozous mauritanus

Predation of the millet head miner was confirmed by molecular analysis of feces or direct observations

Natural pest control by arthropods and birds

19




Grain losses were reduced when panicles were accessible to arthropods only, and even more when they were accessible to both arthropods and birds (-35%).



Effect of trees on insectivorous birds

20


The abundance of insectivorous birds was positively correlated to the diversity of trees at a local scale (SHDI_100 m).



Effect of trees on insectivorous birds

21

The abundance of village weavers was positively correlated to the diversity of trees




Village weavers

Effect of trees on insectivorous birds

22

The abundance of the grey-headed sparrow was negatively correlated to the number of trees at a local scale.



Grey-headed sparrow

Effect of trees on insectivorous bats

23


Palm trees (*B. aethiopum*) serve as a specific nesting site for Taphozous bats.



Effect of trees on insectivorous birds


24

At a landscape scale, no effect of trees on the abundance of insectivorous birds.



Effect of trees on pest incidence

25




Positive effect of baobab trees on egg infestation of panicles but negative effect of neems and balanites. No significant relation between trees and grain loss.

Take-home message

26

- Vertebrates, including birds and bats, are effective predators of the millet head miner.
- They contribute to natural pest control and reduction of millet grain loss.
- The abundance of insectivorous village weaver birds increased with tree diversity whereas that of grey-headed sparrows decreased with the abundance of trees, at a local scale.
- The contribution of trees for natural pest control is species-dependent. Trees can serve as refuges for pest (eg. baobab/moths) and predators (eg. palm tree/bats or neem/village weavers).





Thank you for your attention !

8—10 october 2019

Hotel Ngor Diarama
Dakar, Sénégal

Session 1.2.
Agroforestry

